

Conférence au CRNL le vendredi 30 août 2019



Systems Biology of Mammalian Sleep/Wake Cycles Phosphorylation Hypothesis of Sleep

Hiroki R. UEDA ¹⁻³

1. WPI-IRCIN, UTIAS, The University of Tokyo

2. Systems Pharmacology, Graduate School of Medicine, University of Tokyo

3. Laboratory for Synthetic Biology, Center for Biosystems Dynamics Research, RIKEN

The detailed molecular and cellular mechanisms underlying NREM sleep (slow-wave sleep) and REM sleep (paradoxical sleep) in mammals are still elusive. To address these challenges, we first constructed a mathematical model, Averaged Neuron Model (AN Model), which recapitulates the electrophysiological characteristics of the slow-wave sleep. Comprehensive bifurcation analysis predicted that a Ca^{2+} -dependent hyperpolarization pathway may play a role in slow-wave sleep. To experimentally validate this prediction, we generate and analyze 26 KO mice, and found that impaired Ca^{2+} -dependent K^+ channels (*Kcnn2* and *Kcnn3*), voltage-gated Ca^{2+} channels (*Cacna1g* and *Cacna1h*), or Ca^{2+} /calmodulin-dependent kinases (*Camk2a* and *Camk2b*) decrease sleep duration, while impaired plasma membrane Ca^{2+} ATPase (*Atp2b3*) increases sleep duration. Genetical (*Nr3a*) and pharmacological intervention (PCP, MK-801 for *Nr1/Nr2b*) and whole-brain imaging validated that impaired NMDA receptors reduce sleep duration and directly increase the excitability of cells. Based on these results, we propose **phosphorylation hypothesis of sleep** that phosphorylation-dependent regulation of Ca^{2+} -dependent hyperpolarization pathway underlies the regulation of sleep duration in mammals. We also recently developed a simplified mathematical model, Simplified Averaged Neuron Model (SAN Model), which uncover the important role of K^+ leak channels in NREM sleep. In this talk, I will also describe how we identify essential genes (*Chrm1* and *Chrm3*) in REM sleep regulation, and propose a plausible molecular definition of a paradoxical state of REM sleep.

References

1. Tatsuki et al. *Neuron*, 90(1) : 70–85 (2016).
2. Sunagawa et al, *Cell Reports*, 14(3):662-77 (2016).
3. Susaki et al. *Cell*, 157(3): 726–39, (2014).
4. Tainaka et al. *Cell*, 159(6):911-24(2014).
5. Susaki et al. *Nature Protocols*, 10(11):1709-27(2015).
6. Susaki and Ueda. *Cell Chemical Biology*, 23(1):137-57 (2016).
7. Tainaka et al. *Ann. Rev. of Cell and Devel. Biol.* 32: 713-741 (2016).
8. Ode et al. *Mol. Cell*, 65, 176–190 (2017).
9. Tatsuki et al, *Neurosci. Res.* 118, 48-55 (2017)

10. Ode et al, *Curr. Opin. Neurobiol.* 44, 212-221 (2017)
11. Susaki et al, *NPJ. Syst. Biol. Appl.* 3, 15 (2017)
12. Shinohara et al, *Mol. Cell* 67, 783-798 (2017)
13. Ukai et al, *Nat. Protoc.* 12, 2513-2530 (2017)
14. Shi and Ueda, *BioEssays* 40, 1700105 (2018)
15. Yoshida et al, *PNAS* 115, E9459-E9468 (2018)
16. Niwa et al, *Cell report*, 24, 2231-2247. e7 (2018)